

Legionella Native Valve Endocarditis Case Report and Review of Literature

CASE REPORT

Verena Baumgartner¹,
Gerhard Eich²,
Stefan Christen³,
Corinne Chmiel⁴

- 1 Division of Internal Medicine, City Hospital Waid, Zurich, Switzerland.
- 2 Division of Infectiology, City Hospital Triemli, Zurich, Switzerland.
- 3 Division of Cardiology, City Hospital Waid, Zurich, Switzerland.
- 4 Division of Internal Medicine, City Hospital Waid, Zurich, Switzerland and University Hospital Zurich, Institute of Primary Care, University of Zurich, Switzerland.

Abstract

Background: Legionella native valve endocarditis is exceedingly rare, and to the best of our knowledge, this is the first case complicated by a septic spleen infarction.

Keywords

Legionella; Endocarditis; Pneumonia; Septic Embolism.

Introduction

The incidence of infectious endocarditis worldwide is estimated at 30/100.000 cases per year. In Western Europe and North America endocarditis on native valves show an incidence of 2-6/100.000 per year [1, 2]. Streptococcus and Staphylococcus species are the cause for endocarditis in about 80%, another 10% are caused by Enterococcus species or bacteria of the HACEK-group. According to Barnes et al. culture negative endocarditis accounts for 2.5 to 31% of all cases [3]. Negative blood cultures often result from previous antimicrobial therapy or fastidious bacteria (mostly *Coxiella burnetii*, *Bartonella* spp. And *Tropheryma whippeli*).

Legionella species are very rarely found as causative agent of infectious endocarditis; only 2 patients with involvement of a native valve are mentioned in literature so far. Legionella species are fastidious gram-negative bacilli. Today 57 species are known incorporating at least 79 different serotypes. According to Murder and Yu 20 different Legionella species are considered as pathogens for humans, of which Legionella pneumophila is responsible for 90% of infections [4]. Legionella pneumophila has 16 serotypes, with serotype 1 being the most significant, other Legionella serotypes are found in nosocomial infections. Legionella mostly causes pneumonia by direct

Contact information:

Corinne Chmiel.

PD Dr. med.

Address: University Hospital Zürich,
Institute of Primary Care, University of
Zurich, Pestalozzistrasse 24, CH-8091
Zürich, Switzerland.

Tel: 0041 44 255 98 55.

Fax: 0041 44 255 90 97.

✉ corinne.chmiel@usz.ch


inhalation of contaminated aerosol [5]. Extra pulmonary manifestations are rare and described predominantly in surgical patients as prosthetic valve endocarditis, joint-prosthesis- or wound-infections [6], where the infection pathway is via inoculation with the bacteria during surgery. Gold standard for diagnosis of Legionella pneumonia is a positive culture on a Legionella- BCYE-Agar or a Legionella specific PCR. Since these diagnostic methods are complex and not always available, the most commonly performed testing is the detection of Legionella antigen in the urine, even though the test is only able to detect Legionella pneumophila serotype 1. Serological testing is not useful for diagnosing acute infection since seroconversion often occurs only after several weeks. Final proof of a Legionella endocarditis is a positive histology with typical bacteria seen on electronic microscopy or a positive PCR for Legionella.

Methods and Findings

Case report and review of literature

A 64 year old man with alcoholic liver cirrhosis (Child A) presented with a few weeks' history of fever, dyspnea, cough and thoracic pain. His previous medical history was uneventful and medication intake was reported. On admission he was febrile (40.0°C), had a heart rate of 115/min and a blood pressure of 120/80 mmHg. His respiratory rate was 35/min, the oxygen saturation breathing room air was 91%. The pulmonary examination revealed diminished lung sounds and crackles in the right lower lobe. Heart sounds were normal and there were no murmurs. The electrocardiogram showed a new atrial fibrillation. Leucocytes were at 9.5 G/l (normal range 3.5-10.0 G/l) with a marked left shift (36%), C-reactive protein (CRP) was highly elevated at 333 mg/l (normal range < 5 mg/l), and serum creatinine was 117 µmol/l (normal range 62-106 µmol/l). Urinary spot analy-

Figure 1: Transesophageal echocardiography showing thrombotic material on the aortic valve.



LA: left atrium, LVOT: left ventricular outlet tract, AO: Aorta, AV: Aortic valve.

sis showed microhematuria >10mg/l (normal range <0.3mg/l) and proteinuria of 1g/l (normal range <0.1g/l). An extensive infiltration of the entire right lung was seen on chest X-ray. Community acquired pneumonia was diagnosed and empiric antibiotic treatment with ceftriaxone and clarithromycin initiated. After five days the patient's condition significantly improved and clarithromycin was stopped. After a transient improvement, the patient's inflammatory parameters again deteriorated, and he developed respiratory failure accompanied by delirium tremens. A right basal lung abscess and two fresh spleen infarctions were diagnosed in the thoraco-abdominal CT scan. Transthoracic echocardiography revealed a 6 mm floating thrombus on the aortic valve, which was confirmed by transesophageal echocardiography (**Figure 1**). According to the Duke criteria (1 major, 4 minor criteria) diagnosis of culture negative endocarditis with suspicion of septic embolisms was made. The urinary test result for Legionella antigen was positive and antibiotic therapy with clarithromycin was restarted. Consecutively, a Legionella antibody analysis of the serum by complement binding reaction was performed showing a slightly positive result. Repetitive blood cultures, specific PCR

for Legionella as well as conventional culture of the pleural fluid remained negative, as did Brucella serology. Consecutively, the patient's pulmonary and cardiac status steadily improved. Twenty days after admission the patient was transferred to an alcohol rehabilitation institution. After a total of six weeks antibiotic treatment with clarithromycin the patient was seen for follow-up. He presented in good clinical condition with normalization of the chest X-ray and laboratory parameters. The repeated analysis of Legionella specific PCR and broad spectrum PCR of the serum were negative. Legionella urinary antigen analysis remained positive. Although transthoracic echocardiography still showed some valve abnormalities interpreted as residual scars, we decided to stop the antibiotic treatment. At follow up three months later the patient was well and asymptomatic.

Discussion

The current case is, to the best of our knowledge, the first report of an endocarditis due to Legionella complicated by a spleen abscess. Altogether, it represents the third report of a native valve endocarditis attributable to Legionella pneumophila (Table 1).

Legionella endocarditis has rarely been described in literature. Only 16 cases have been reported from which the first is dated to 1984 [7]. In almost all cases patients had experienced valve surgery, only two other cases besides ours had native valves [8, 9]. In our case the urinary and serum analyses for Legionella antigen was positive, but specific PCR for Legionella as well as was broad spectrum PCR showed repeatedly negative results. In literature only in one patient the diagnosis was confirmed with PCR and sequencing from aortic tissue [9], in the other, Legionella pneumophila serotype 1 was detected in the urine sample and cultures on buffered charcoal-yeast extract (BCYE)-Agar using blood and bronchial fluid [8].

Admission diagnosis of our patient was pneumonia and antibiotic treatment with Ceftriaxone and Clarithromycin was started. Since ceasing the therapy with Clarithromycin was followed by serious deterioration of the patient's state, we assumed systemic spreading of the bacteria. Diagnosis of endocarditis was established by Duke criteria fulfilling 1 major criteria by positive transesophageal echocardiography (Figure 1) with a floating mass on the aortic valve, as well as 4 minor criteria: fever (temperature > 38°C), vascular phenomena (lung and spleen infarction), immunological phenomena (nephritic syndrome) and microbiological evidence (positive urinary antigen for Legionella pneumophila serotype 1 and positive blood serology).

Embolic events due to Legionella endocarditis seem to be very rare and have been reported in only two other cases with valve-prosthesis infection [10, 11]. Our patient showed two fresh spleen infarctions diagnosed in thoraco-abdominal CT scan. Whether they were caused by septic or thrombotic cardiac embolism triggered by the paroxysmal atrial fibrillation is not certain, the lack of other thrombotic material in transthoracic and transesophageal echocardiography other than the floating vegetation on the aortic valve as well as the presence of immunological phenomena (e.g. nephritic syndrome) seem to favor septic embolism as a cause.

Regarding the duration of therapy for Legionella endocarditis evidently no guidelines exists. Therapy regimens ranging from six weeks for native valves [8] to 5-14 months for prosthetic valves [10, 12] have been described in literature (Table 1).

Conclusion

In case of culture negative endocarditis Legionella species has to be thought of as possible causative agent, particularly in patients with prosthetic valves. Embolic events in Legionella endocarditis are extremely rare, but have been reported. Molecular proof of the bacteria with PCR respectively silver

Table 1. Literature Review Legionella Endocarditis (all Case Reports).

Case	Author	Year	Valve type	Diagnosis	Patient characteristics/ risk factors	Treatment	Complications/ outcome	Pathogen
1	McCabe R E et al.	1984	pr, po, a, m	Valve culture, serology	W, 60 y, rheumatic valvular disease	ERM for 6 mo, RIF for 2 mo - valve replacement	Cured	L. Pneumophila
2	Tompkins L S et al.	1988	pr, po, a, m	Valve culture, serology	Valve surgery at sumc during a general outbreak of l.	ERM 4g iv for 8 wk, 2g po for 12 mo, RIF 600mg po for 6 mo - valve replacement	Cured at 48 mo follow up	L. Pneumophila
3	Tompkins L S et al.	1988	pr, po, a	Valve culture, serology	Valve surgery at sumc during a general outbreak of l.	ERM 4g iv for 6 mo, RIF 600mg po for 6 mo, vancomycin 2 g iv for 6 mo - valve replacement	Cured at 40 mo follow up	L. Dumoffii
4	Tompkins L S et al.	1988	pr, me	Serology	Valve surgery at sumc during a general outbreak of l.	ERM 4g iv for 2 mo, 2g po for 6 mo, RIF 600mg po for 6 mo	Cured	L. Pneumophila
5	Tompkins L S et al.	1988	pr, po, a	Valve culture, serology, blood culture	Valve surgery at sumc during a general outbreak of l.	ERM 2g iv for 9 mo, RIF 1200mg po 9 mo- valve replacement	Cured at 11 mo follow up	L. Dumoffii
6	Tompkins L S et al.	1988	pr, po, m	Valve culture, serology, blood culture	Valve surgery at sumc during a general outbreak of l.	ERM 3g iv for 5wk, followed by ERM 4g iv for 1 mo, followed by 2g po and RIF 600mg po for 14 mo - valve replacement	Cured	L. Dumoffii
7	Tompkins L S et al.	1988	pr, po, m	Blood culture, serology	Valve surgery at sumc during a general outbreak of l.	ERM 4g iv for 2 days, followed by Ciprofloxacin 600mg iv and RIF 1200mg po for 10 wk - valve replacement	Development of mitral insufficiency	L. Pneumophila
8	Tompkins L S et al.	1988	pr, po	Blood culture, serology	Valve surgery at sumc during a general outbreak of l.	ERM 4g iv for 2 mo plus RIF 1200 mg po for 5 mo	Cured	L. Pneumophila, l. Dumoffii
9	Park D et al.	1994	pr, me, a	Serology	MI, 65 y, diabetic	Doxycyclin - valve replacement	Microscopic hematuria	L. Micdadei
10	Chen TT et al.	1996	pr, me, a	Valve culture, blood culture	MI, 33 y, marfan syndrome	ERM for 6 mo, RIF - valve replacement	Pseudoaneurysm	L. Pneumophila

Case	Author	Year	Valve type	Diagnosis	Patient characteristics/risk factors	Treatment	Complications/outcome	Pathogen
11	R. Massey et al.	2003	pr, h, a	Valve cultures negative after prolonged antibiotic treatment, I. Pneumonia microbiologically confirmed (method not further specified)	MI, 26 y, congenital heart disease, I. Pneumonia during a localized outbreak	Clarithromycin, RIF iv - repeat valve replacement	Microembolism of digits, respiratory failure, dic	L. Pneumophila
12	Patel MC et al.	2005	pr, b, a	Valve culture	MI, 63 y, kidney transplantation, history of hodgkin's disease	Levofloxacin indefinitely - valve replacement	No relapse	L. Micdadei
13	Samuel V et al.	2011	native	Blood culture, bal, I. Antigen urine test	W, 42 y, corticosteroids for pneumonitis	Levofloxacin for 6 wk, RIF for 6 wk	Pneumonia, ards, dic, septic shock, acute kidney injury with hemodialysis, cured	L. Pneumophila
14	M. M. Pearce et al.	2011	native	Valve culture	W, 68 y, good pasture syndrome, kidney transplantation, cmv infection	(Penicillin allergy) Vancomycin, Ciprofloxacin, Gentamycin - valve replacement - after positive culture Moxifloxacin for 6 wk - relapse: vancomycin, RIF Azithromycin, valve replacement was refused	Death	New I. Strain, designated h63, shares the highest similarity with I. Brunensis
15	Yuriko Fukuta et al.	2012	pr, me, m	Valve culture	W, 57 y, sle and anti-phospholipid-sydnrom, corticosteroids, azathioprin	Levofloxacin for 5 mo - valve replacement	Brain abscess, after 5 mo of therapy persistence of frontal brain mass, therapy was continued (duration unknown)	L. Micdadei
16	our case	2013	native	L. Antigen urine test, serology	MI, 64 y, alcoholic liver cirrhosis, I. Pneumonia	Clarithromycin for 6 wk	Spleen infarction, cured 90 mo follow up	L. Pneumophila

Prosthetic (pr), porcine (po), aortic (a), mitral (m), mechanical (me), homograft (h), bovine (b), stanford university medical center (sumc), cytomegaic virus (cmv), systemic lupus erythematodes (sle), rifampicin, legionella (l), women (f), male (ml), years (y), cytmoeagal virus (cmc), broncho alveolar lavage (bal), inveravenous (iv), peroral (po), months (mo), weeks (wk), disseminated intravascular coagulopathy (dic)

impregnation stains from histological material are desirable, but unfortunately histological material is not always available.

Potential conflicts of interest

All authors declare no conflict of interests. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest.

References

1. Hoge vik H, et al. Epidemiologic aspects of infective endocarditis in an urban population. A five-year prospective study. *Medicine* 1995; 74: 324-39.
2. Berlin JA, Abrutyn E, Strom BL, Kinman JL, Levison ME, Korzeniowski OM, et al. Incidence of infective endocarditis in the Delaware Valley, 1988-1990.
3. Barnes PD, Crook DWM. Culture negative endocarditis. *J Infect* 1997; 35: 209-13.
4. Robert R. Muder and L. Yu Victor. Infection Due to Legionella Species Other Than *L. pneumophila*. *Clin Infect Dis*. (2002) 35 (8): 990-998
5. Carratalà J, Garcia-Vidal C. An update on Legionella. *Curr Opin Infect Dis* 2010; Apr; 23(2): 152-7.
6. Lowry PW, Tompkins LS. Nosocomial legionellosis: a review of pulmonary and extrapulmonary syndromes. 1993 Feb; 21(1): 21-7.
7. Prendergast BD. Diagnosis of infective endocarditis. *BMJ*. 2002 Oct 19; 325(7369): 845-6.
8. Samuel V., Bajwa A. A., Cury J. D. First case of Legionella pneumophila native valve endocarditis. *Int J Infect Dis*. 2011 Aug; 15(8): e576-7.
9. N. Theodoropoulos, G. A. Noskin, J. P. Flaherty. Native Valve Endocarditis Due to a Novel Strain of Legionella. *J Clin Microbiol*. 2011 September; 49(9): 3340-3342.
10. Yuriko Fukuta, Isil Z. Yildiz-Aktas, A. William Pasculle & Peter J. Veldkamp. Legionella micdadei prosthetic valve endocarditis complicated by brain abscess: Case report and review of the literature. *Scandinavian Journal of Infectious Diseases*, 2012; 44: 414-418.
11. R. Massey, P. Kumar, J. Pepper. Innocent victim of a localized outbreak: legionella endocarditis. *Heart*. 2003 May; 89(5): e16.
12. Brouqui P, Raoult D. Endocarditis due to rare and fastidious bacteria. *Clin Microbiol Rev* 2001; 14: 177-207.

Publish in International Archives of Medicine

International Archives of Medicine is an open access journal publishing articles encompassing all aspects of medical science and clinical practice. IAM is considered a megajournal with independent sections on all areas of medicine. IAM is a really international journal with authors and board members from all around the world. The journal is widely indexed and classified Q1 in category Medicine.